



Diplomate, American Board of Endodontics

12791 Newport Avenue, Suite 102 | Tustin, California 92780 | T: 714.838.1415 | F: 714.838.2537 | Info@Pacific-Endo.com

## **Post-Procedure Instructions**

Your root canal procedure has been **completed / initiated**:

- Your tooth and gum tissues will be sore for several days due to inflammation—this is normal and consistent with healing. Inflammation increases and peaks 72 hours after your appointment, then gets better.
  - You may feel soreness around the tooth, the gum area where the shots were given.
    Rinsing with warm salt water may provide relief.
  - There may be some tenderness upon biting. This is also normal. Try to avoid biting on the tooth worked on to avoid additional inflammation.
  - Lastly, your jaw muscles may be sore on both sides as a result of keeping your mouth open for an extended period of time. If this happens, stretch the muscles or chew gum on the side not worked on.

•	The following medication(5) is are devised around the clock for the next 40 12 hours for
	pain/inflammation:
	☐ Ibuprofen (Advil/Motrin) 400-600mg, every 4-6 hours, up to 3200mg per day. If ibuprofen
	alone is not alleviating your discomfort, you may supplement with the acetaminophen
	regimen below. Take Ibuprofen with food.
	☐ Acetaminophen (Tylenol) 500-1000mg, every 4-6 hours, up to 3000mg per day.
•	The following antibiotic is advised; please take to completion:
	☐ Amoxicillin
	☐ Azithromycin/Z-Pak
•	Additionally:
<b>~</b> .,	continue to be numb for a few bourse the numbered will gradually wear off While you are still

The following medication(s) is/are advised around the clock for the next 48-72 hours for

You may **continue to be numb for a few hours**; the numbness will gradually wear off. While you are still numb, a soft mechanical diet (e.g. soups, smoothies, shakes, etc.) is advised to limit potentially biting yourself.

Your tooth has been filled with a temporary filling. Please allow at least one hour for the temporary filling to set before eating anything. Even so, avoid eating on the tooth as much as possible. You may feel like the filling has fallen out when in all likelihood it has been pushed further into the tooth. It is imperative that you schedule a permanent restoration with your dentist as soon as possible.

Should you have additional questions or concerns, please call our office at (714)838-1415.